


## MCTs - BENEFITS BEYOND THE BRAIN

MCTs (Medium-chain triglycerides) are a unique fat source with vast proven benefits, with powerful potential extending beyond just neurological health.



### REFERENCES

- McCartt, M. F., & DiNicolantonio, J. J. (2016). Lauric acid-rich medium-chain triglycerides can substitute for other oils in cooking applications and may have limited pathogenicity. *Open Heart*, 3(2), e000467.
- Leung, Y. B., Cave, N. J., Heiser, A., Edwards, P. J. B., Godfrey, A. J. R., & Wester, T. (2020). Metabolic and immunological effects of intermittent fasting on a ketogenic diet containing medium-chain triglycerides in healthy dogs. *Frontiers in Veterinary Science*, 480.
- Taha, A. Y., Henderson, S. T., & Burnham, W. M. (2009). Dietary enrichment with medium chain triglycerides (AC-1203) elevates polyunsaturated fatty acids in the parietal cortex of aged dogs: implications for treating age-related cognitive decline. *Neurochemical Research*, 34, 1619–1625.
- Pan, Y., Larson, B., Araujo, J. A., Lau, W., de Rivera, C., Santana, R., Gore, A., & Milgram, N. W. (2010). Dietary supplementation with medium-chain TAG has long-lasting cognition-enhancing effects in aged dogs. *The British Journal of Nutrition*, 103(12), 1746–1754.
- Law, T. H., Volk, H. A., Pan, Y., Zanghi, B., & Want, E. J. (2018). Metabolic perturbations associated with the consumption of a ketogenic medium-chain TAG diet in dogs with idiopathic epilepsy. *British Journal of Nutrition*, 120(5), 484–490.
- Molina, J., Jean-Philippe, C., Conboy, L., Añor, S., de la Fuente, C., Wrzosek, M. A., Spycher, A., Luchsinger, E., Wenger-Rüegg, B., & Montoliu, P. (2020). Efficacy of medium chain triglyceride oil dietary supplementation in reducing seizure frequency in dogs with idiopathic epilepsy without cluster seizures: a non-blinded, prospective clinical trial. *Veterinary Record*, 187(9), 356.
- Kathrani, A., & Parkes, G. (2022). A Preliminary Study of Modulen IBD Liquid Diet in Hospitalized Dogs with Protein-Losing Enteropathy. *Animals*, 12(12), 1594.
- Shah, N. D., & Limketkai, B. N. (2017). The use of medium-chain triglycerides in gastrointestinal disorders. *Practical Gastroenterology*, 160, 20–25.
- Vecchiato, C. G., Pinna, C., Sung, C. H., Borrelli De Andreis, F., Suchodolski, J. S., Pilla, R., Delsante, C., Sportelli, F., Mammi, L. M. E., Pietra, M., & Biagi, G. (2023). Fecal Microbiota, Bile Acids, Sterols, and Fatty Acids in Dogs with Chronic Enteropathy Fed a Home-Cooked Diet Supplemented with Coconut Oil. *Animals*, 13(3), 502.
- Grancher, D., Jean-Blain, C., Frey, A., Schirardin, H., & Bach, A. C. (1987). Studies on the tolerance of medium chain triglycerides in dogs. *Journal of parenteral and enteral nutrition*, 11(3), 280–286.
- Wilshaw, J., Boswood, A., Chang, Y. M., Sands, C. J., Camuzeaux, S., Lewis, M. R., Xia, D., & Connolly, D. J. (2022). Evidence of altered fatty acid metabolism in dogs with naturally occurring valvular heart disease and congestive heart failure. *Metabolomics*, 18(6), 34.
- Li, Q., Heaney, A., Langenfeld-McCoy, N., Boler, B. V., & Laflamme, D. P. (2019). Dietary intervention reduces left atrial enlargement in dogs with early preclinical myxomatous mitral valve disease: a blinded randomized controlled study in 36 dogs. *BMC Veterinary Research*, 15(1), 425.
- Li, Q. (2022). Metabolic reprogramming, gut dysbiosis, and nutrition intervention in canine heart disease. *Frontiers in Veterinary Science*, 9, 791754.
- Hall, J. A., & Jewell, D. E. (2012). Feeding healthy beagles medium-chain triglycerides, fish oil, and carnitine offsets age-related changes in serum fatty acids and carnitine metabolites. *PLoS One*, 7(11), e49510.